

# Post Graduate School Indian Agricultural Research Institute, New Delhi

# Examination for Admission to Ph.D. Programme 2013-2014

Discipline	: Genetics					
Discipline Code	: 12	Roll No.				

### **Please Note:**

- (i) This question paper contains 13 pages. Please check whether all the pages are printed in this set. Report discrepancy, if any, immediately to the invigilator.
- (ii) There shall be NEGATIVE marking for WRONG answers in the Multiple Choice type questions (No. 1 to 130) which carry one mark each. For every wrong answer 0.25 mark will be deducted.

## PART – I (General Agriculture)

Multiple choice questions (No. 1 to 30). Choose the correct answer (a, b, c or d) and enter your choice in the circle (by shading with a pencil) on the OMR - answer sheet as per the instructions given on the answer sheet.

- Who is the present Chairman of Protection of Plant Varieties and Farmers' Right Authority (PPV&FRA)?
- a) Dr. R.R. Hanchinal
- b) Dr. P.L. Gautam
- c) Dr. S. Nagarajan
- d) Dr. Swapan K. Datta
- 2. Which among the following is another name for vitamin B<sub>12</sub>?
- a) Niacin
- b) Pyridoxal phosphate
- c) Cobalamin
- d) Riboflavin
- 3. The largest share in India's farm export earning in the year 2011-12 was from
- a) Basmati rice
- b) Non-basmati rice
- c) Sugar
- d) Guar gum
- 4. The National Bureau of Agriculturally Important Insects was established by ICAR in \_\_\_\_\_\_, was earlier known as \_\_\_\_\_.
- a) Bangalore; PDBC
- b) New Delhi; National Pusa Collection
- c) Ranchi; Indian Lac Research Institute
- d) New Delhi; NCIPM

- 5. The most important sucking pests of cotton and rice are respectively
- a) Nilaparvata lugens and Aphis gossypii
- b) Aphis gossypii and Thrips oryzae
- c) Amrasca biguttula biguttula and Scirtothrips dorsalis
- d) Thrips gossypii and Orseolia oryzae
- 6. Which of the following microorganism causes fatal poisoning in canned fruits and vegetables?
- a) Aspergillus flavus
- b) Penicillium digitatum
- c) Clostridium botulinum
- d) Rhizoctonia solani
- 7. The cause of the great Bengal Famine was
- a) Blast of rice
- b) Brown spot of rice
- c) Rust of wheat
- d) Karnal bunt of wheat
- 8. Actinomycetes belong to
- a) The fungi
- b) Eukaryote
- c) Mycelia sterilia
- d) None of the above
- 9. A virus-free clone from a virus infected plant can be obtained by
- a) Cotyledonary leaf culture
- b) Axenic culture
- c) Stem culture
- d) Meristem tip culture
- 10. Which of the following is not an objective of the National Food Security Mission?
- Sustainable increase in production of rice, wheat and pulses
- Restoring soil fertility and productivity at individual farm level
- Promoting use of bio-pesticides and organic fertilizers
- d) Creation of employment opportunities

- Agmarknet, a portal for the dissemination of agricultural marketing information, is a joint endeavour of
- a) DMI and NIC
- b) DMI and Ministry of Agriculture
- c) NIC and Ministry of Agriculture
- d) DMI and Directorate of Economics and Statistics
- The share of agriculture and allied activities in India's GDP at constant prices in 2011-12 was
- a) 14.1%
- b) 14.7%
- c) 15.6%
- d) 17.0%
- 13. The average size of land holding in India according to Agricultural Census 2005-06 is
- a) 0.38 ha
- b) 1.23 ha
- c) 1.49 ha
- d) 1.70 ha
- 14. 'Farmers First' concept was proposed by
- a) Paul Leagans
- b) Neils Rolling
- c) Robert Chamber
- d) Indira Gandhi
- 15. In the year 2012, GM crops were cultivated in an area of
- a) 150 million hectare in 18 countries
- b) 170 million hectare in 28 countries
- c) 200 million hectare in 18 countries
- d) 1.70 million hectare in 28 countries
- The broad-spectrum systematic herbicide glyphosate kills the weeds by inhibiting the biosynthesis of
- a) Phenylalanine
- b) Alanine
- c) Glutamine
- d) Cysteine
- 17. At harvest, the above ground straw (leaf, sheath and stem) weight and grain weight of paddy crop are 5.5 and 4.5 tonnes per hectare, respectively. What is the harvest index of paddy?
- a) 45%
- b) 50%
- c) 55%
- d) 100%
- Crossing over between non-sister chromatids of homologous chromosomes takes place during
- a) Leptotene
- b) Pachytene
- c) Diplotene
- d) Zygotene

- 19. The term 'Heterosis' was coined by
- a) G.H. Shull
- b) W. Bateson
- c) T.H. Morgan
- d) E.M. East
- 20. When a transgenic plant is crossed with a non-transgenic, what would be the zygosity status of the F<sub>1</sub> plant?
- a) Homozygous
- b) Heterozygous
- c) Hemizygous
- d) Nullizygous
- 21. The highest per capita consumption of flowers in the world is in
- a) The USA
- b) India
- c) Switzerland
- d) The Netherlands
- 22. Which of the following is a very rich source of betalain pigment?
- a) Radish
- b) Beet root
- c) Carrot
- d) Red cabbage
- 23. Dog ridge is
- a) Salt tolerant rootstocks of mango
- b) Salt tolerant rootstocks of guava
- c) Salt tolerant rootstocks of grape
- d) Salt tolerant rootstocks of citrus
- 24. Which of the following micronutrients are most widely deficient in Indian soils?
- a) Zinc and boron
- b) Zinc and iron
- c) Zinc and manganese
- d) Zinc and copper
- 25. Which of the following fertilizers is not produced in India?
- a) DAP
- b) Urea
- c) Muriate of potash
- d) TSP
- 26. What is the estimated extent of salt affected soils in India?
- a) 5.42 mha
- b) 7.42 mha
- c) 11.42 mha
- d) 17.42 mha
- 27. Which of the following is not a feature of watershed?
- a) Hydrological unit
- b) Biophysical unit
- c) Socio-economic unit
- d) Production unit

- 28. Correlation coefficient 'r' lies between
- a) 0 and 1
- b) -1 and 1
- c) -1 and 0
- d) 0 and  $\infty$
- 29. For the data 1, -2, 4, geometric mean is
- a) 2
- b) 4
- d) -2
- 30. The relationship between Arithmetic mean (A), Harmonic mean (H) and Geometric mean (G) is
- a)  $G^2=AH$
- b)  $G=\sqrt{A+H}$ c)  $H^2=GA$
- d)  $A^2 = GH$

# PART - II (Subject Paper)

Multiple choice questions (No. 31 to 130). Choose the correct answer (a, b, c or d) and enter your choice in the circle (by shading with a pencil) on the OMR answer sheet as per the instructions given on the answer sheet.

- 31. If genes A and B are linked and separated by 14 map units, the percentage of parental progeny produced from test crossing a double heterozygote will be
- a) 7
- b) 14
- c) 46
- d) 86
- 32. The shortest phase of the cell cycle is
- a) G<sub>2</sub>
- b) G<sub>1</sub>
- Μ c)
- d) S
- 33. When the co-efficient of coincidence is greater than 1, then
- a) Expected double crossovers are less than observed double crossovers
- b) Expected double crossovers are more than observed double crossovers
- c) Interference is 1
- d) Interference is 0
- 34. Genes that are greater than 50 map units apart cannot be used to predict the frequency of recombinant products, because
- a) Of the absence of recombination
- b) The genes will segregate independently
- c) Of excess of single crossovers
- d) Coefficient of coincidence will be 1

- 35. A paracentric inversion involves chromosomal
- a) Region including the centromere
- b) Region excluding the centromere
- Exchange between inverted centromeric regions of 2 chromosomes
- Exchange between telomeric regions of 2 chromosomes
- 36. If 'n' denotes the number of alleles of a gene, the number of genotypes in a multiple allelic series is given by
- n(n-1)
- b) n(n-1)
- d) n(n+1)
- 37. X/A ratio in a male Drosophila melanogaster will be
- 1.5
- b) 1.0
- 0.75 c)
- d) 0.5
- 38. A+G / T+C ratio in one strand of DNA is 0.8, what will be this ratio in the complementary strand?
- 0.20 a)
- b) 0.80
- 1.60 c)
- d) 1.25
- 39. In bread wheat (T. aestivum L.), maximum number of multivalents will be observed in stock.
- Nulli-5A
- b) Nulli-5B
- c) Nulli-5D
- d) Nulli-3B
- 40. Which of the following is not a method of population improvement?
- Pure line selection
- b) Mass selection
- Reciprocal recurrent selection
- Diallel selective mating system
- 41. Gene whose expression is specific to the time and type of cell is called
- a) Modifier gene
- b) Operator gene
- Homeotic gene c)
- d) Temporal gene

- 42. In Lac operon, \_\_\_\_\_ acts as an inducer.
- a) Lactose
- b) Allolactose
- c) Glucose
- d) Galactose
- 43. Number of different types of secondary trisomics possible in maize is
- a) 10
- b) 20
- c) 30
- d) 40
- 44. Which of the following can alter the arm ratio of a chromosome without changing its length?
- a) Deletion
- b) Paracentric inversion
- c) Pericentric inversion
- d) Duplication
- 45. How many genetically different gametes can be produced by an individual of genotype AaBbccDDEe?
- a) 5
- b) 8
- c) 10
- d) 32
- 46. Base pair substitutions involving replacement of a purine for a pyrimidine base and *vice versa* are called
- a) Transition
- b) Frameshift
- c) Transcription
- d) Transversion
- 47. The bases in DNA are stacked at a spacing of
- a) 3.4 A°
- b) 34 A°
- c) 0.34 A°
- d) 0.034 A°
- 48. A double tetrasomic is represented by the notation
- a) 2n+2
- b) 2n+2+2
- c) 2n-2
- d) 2n-1
- 49. With 20 inbred lines, the number of possible three way crosses (excluding reciprocals) would be
- a) 190
- b) 400
- c) 3420
- d) 14535

- Balanced polymorphism in nature occurs only due to
- a) Homozygous predominance
- b) Heterozygous advantage
- c) Elimination of recessives
- d) Complete dominance
- 51. Overlapping genes were first discovered in
- a) E. coli
- b) Neurospora
- c) Bacteriophage
- d) Arabidopsis
- 52. SNP refers to
- a) Sequenced nucleotides polymorphism
- b) Super nucleotide polymorphism
- c) Single nucleotide polymorphism
- d) Saturated nucleotide polymorphism
- 53. The position of a gene with respect to adjacent region influences its expression. This phenomenon was first demonstrated in
- a) Maize
- b) Pea
- c) Fruitfly
- d) Bacteriophage
- 54. Specific genes which expresses either in specialized cells or respond to specific stimuli are described as
- a) House keeping genes
- b) Smart genes
- c) Producer genes
- d) Integrator genes
- 55. The proportion of heterozygotes at a single locus in  $F_4$  generation of a cross will be
- a) 1/2
- b) 1/4
- c) 1/8
- d) 1/16
- 56. A single chromatid contains
- a) A single DNA double helix
- b) Two DNA double helices
- c) Many DNA double helices
- d) Single strand of DNA
- 57. What will be the proportion of completely homozygous individuals in F<sub>2</sub> of a dihybrid cross?
- a) 1/2
- b) 1/4
- c) 1/8
- d) 1/16
- 58. Which among the following is a 'start' codon in eukaryotes?
- a) AUU
- b) AUC
- c) AUA
- d) AUG

- 59. While calculating linkage in a certain 3-point test cross, the interference was 0.963. It implies that \_\_\_\_\_\_ of the expected double crossovers \_\_\_\_\_.
- a) 9.63%, were not observed
- b) 96.3%, were not observed
- c) 9.63%, were observed
- d) 96.3%, were observed
- 60. Breakage of homologous chromatids and non-reciprocal exchange of its part is referred to as
- a) Recombination
- b) Unequal crossing over
- c) Gene conversion
- d) Translocation
- 61. The term 'genome' was coined by
- a) Bateson
- b) Stebbins
- c) Muller
- d) Winkler
- 62. The first plant disease resistance gene to be cloned and characterized was
- a) Lr gene in wheat
- b) Hm1 gene in maize
- c) Chitinase gene in rice
- d) Rps1 gene in Arabidopsis
- 63. What would be the sequence of bases in the mRNA transcript generated from transcription of this short stretch of DNA:

5' ATTC 3' 3' TAAG 5'

- a) 5' ATTC 3'
- b) 5' TAAG 3'
- c) 5' AUUC 3'
- d) 5' UAAG 3'
- 64. Which of the following is not a feature of eukaryotic mRNA processing?
- a) Alignment of the Shine-Dalgarno sequence
- b) Addition of the 5' methylguanosine cap
- c) Addition of the 3 poly-A tail
- d) Removal of the introns
- 65. The presence-absence hypothesis to explain mutation was proposed by
- a) Hugo deVries
- b) H.J. Muller
- c) W. Bateson
- d) C. Correns
- 66. Proteins are quantified by
- a) Southern blot
- b) Northern blot
- c) Western blot
- d) Eastern blot

- 67. In maize, there are 10 pairs of chromosomes in normal sporophyte tissues. Then, 30, 20 and 10 number of chromosomes are expected in which parts of the plant?
- a) Embryo, root tip, leaf
- b) Embryo sac, seed, root tip
- c) Endosperm, root tip, pollen tube nucleus
- d) Root tip, pollen tube nucleus, leaf
- 68. Coefficient of variation (%) is computed by the formula:

a) 
$$\frac{\text{SE}}{\overline{X}} \times 100$$

b) 
$$\frac{SD}{\sqrt{\overline{X}}} \times 100$$

c) 
$$\frac{SD}{\overline{X}} \times 100$$

d) 
$$\frac{V}{\bar{X}} \times 100$$

- 69. In artificial (manual) hybridization program, the performed activities are (given in random order):
  - i) Pollination, ii) Tagging,
  - iii) Bagging, iv) Emasculation

The correct order of these activities is

- a) iii, iv, i, ii
- b) iv, ii, i, iii
- c) iv, iii, i, ii
- d) ii, iv, i, iii
- 70. Synaptonemal complex is formed in
- a) Zygotene stage
- b) Pachytene stage
- c) Leptotene stage
- d) Diplotene stage
- 71. Under self-fertilization, homozygosity increases in each generation at the rate of
- a) 25%
- b) 50%
- c) 75%
- d) 12.5%
- 72. Selection in favour of both extreme phenotypes is referred to as
- a) Stabilizing selection
- b) Directional selection
- c) Disruptive selection
- d) Cyclic selection
- 73. The Protection of Plant Varieties and Farmers' Rights Act was passed by the Indian Parliament in the year
- a) 1994
- b) 1999
- c) 2001
- d) 2003

- 74. Essentially Derived Varieties (EDVs) are produced by
- a) Mutation
- b) Backcross breeding
- c) Genetic transformation
- d) All of the above
- 75. Inbreeding depression was first observed in maize by
- a) G.H. Shull
- b) K. Mather
- c) N.L. Dhawan
- d) D. Lewis
- 76. Maintenance breeding refers to
- Upgrading resistance to disease in crop variety
- b) Improving nutritive quality of produce
- c) Improving genetic purity of a crop variety
- d) Maintain original yield level of a crop variety
- 77. Which one of the following analysis is appropriate to measure genetic diversity among lines?
- a) Diallel analysis
- b) Line × Tester analysis
- c) D<sup>2</sup> analysis
- d) Metroglyph analysis
- 78. Inbreeding in a population influences
- a) Gene and genotype frequencies
- b) Genotype frequency only
- c) Gene frequency only
- d) Neither gene nor genotype frequency
- 79. Which among the following is genetically heterogeneous but homozygous?
- a) Hybrid
- b) Composite
- c) Multiline
- d) Synthetic
- 80. If a plant is heterozygous for four allelic pairs then upon selfing how many types of genotype are possible?
- a) 9
- b) 27
- c) 64
- d) 81
- 81. If the blood group of the husband is 'AB' and that of the wife is 'O', which among the following lot of siblings with the indicated blood groups are likely to contain their offsprings?
- a) A and B
- b) B and O
- c) AB and O
- d) A and AB

- 82. Which among the following classes of seeds have highest purity?
- a) Breeder's seed
- b) Nucleus seed
- c) Certified seed
- d) TL seed
- 83. In which of the following mapping functions, it is presumed that 'interference' is zero?
- a) Fisher's mapping function
- b) Haldane's mapping function
- c) Kosambi's mapping function
- d) Sturtevant's mapping function
- 84. First linkage map was constructed by
- a) T.H. Morgan
- b) A.H. Sturtevant
- c) S.R. McCouch
- d) B. McClintock
- 85. Heritability of a molecular marker locus is
- a) 0%
- b) 50%
- c) 100%
- d) 150%
- 86. Which among the following markers is best suited for testing hybridity of a plant?
- a) RAPD
- b) SSR
- c) AFLP
- d) ISSR
- 87. If a plant with the genotype AaBbCc is self-fertilized, what proportion of progeny plants will be of the genotype AabbCc (assume that the genes are assorting independently)?
- a) 1/2
- b) 1/8
- c) 1/16
- d) 1/32
- 88. A top cross is a cross between
- a) Two contrasting inbreds
- b) Inbred and an open pollinated variety
- c) An F<sub>1</sub> and an open pollinated variety
- d) A double cross hybrid and an inbred
- 89. ZFN stands for
- a) Zinc finger nitrate
- b) Zinc finger nuclease
- c) Zebra fish number
- d) Zinc finger normal
- 90. Which among the following is a multiparent mapping population?
- a) RIL
- b) NIL
- c) MAGIC
- d) F<sub>2:3</sub>

- 91. The seeds produced on a F<sub>1</sub> plant represent which generation?
- a) F<sub>0</sub>
- b) F<sub>1</sub>
- c) F<sub>2</sub>
- d)  $F_3$
- 92. In a 'cis-trans' test, if two mutations complement each other, then
- a) Both the mutations are structurally allelic
- b) Both the mutations are structurally non-allelic
- Both the mutations are functionally nonallelic
- d) Both b) & c)
- 93. Who among the following geneticists established for the first time that 'gene is sub-divisible'?
- a) Seymour Benzer
- b) Clarence P. Oliver
- c) Edward Lewis
- d) Charles Yanofsky
- 94. Who among the following was not associated in cracking the genetic codes?
- a) Marshall Nirenberg
- b) H.G. Khorana
- c) F.C. Crick
- d) J.D. Watson
- 95. A typical bi-parental cross is performed with an aim to
- a) Decrease deletorious alleles
- b) Decrease homozygosity
- c) Increase frequency of desirable alleles
- d) Increase heterozygosity
- 96. The blue color tag of a seed packet indicates that it is
- a) Breeder's seed
- b) Nucleus seed
- c) Certified seed
- d) TL seed
- 97. The differences in expression of the same gene inherited from the mother and the father is explained by the phenomenon of
- a) Paramutation
- b) Epistasis
- c) Gene silencing
- d) Imprinting
- 98. The polytene chromosomes are thick because they are
- a) Highly condensed
- b) Heavily coated with protein
- c) Multistranded
- d) Rich in heterochromatin

- 99. The cell deprived of Uracil will show
- a) No DNA replication
- b) No protein synthesis
- c) No cell division
- d) Mitotic crossing over
- 100. Cytosine gives rise to Uracil through
- a) Methylation
- b) Demethylation
- c) Oxidative deamination
- d) None of the above
- 101. Teminism refers to
- a) DNA denaturation
- b) RNA denaturation
- c) Reverse transcription
- d) DNA-RNA hybridization
- 102. Baldness in humans is an example of
- a) Sex linked trait
- b) Sex influenced trait
- c) Y-linked trait
- d) Sex limited trait
- 103. If the DNA of a species has the mole fraction of A+T = 0.44, the mole fraction of G (Guanine) will be
- a) 0.22
- b) 0.28
- c) 0.44
- d) 0.56
- 104. Consider a biallelic locus in a population following Hardy-Weinberg equilibrium. If the genotype frequency of the recessive homozygote is 0.04, what would be the allelic frequency of dominant allele?
- a) 0.02
- b) 0.04
- c) 0.2
- d) 0.8
- 105. The bulk method of breeding was first used by
- a) Nilsson-Ehle
- b) Brim
- c) Suneson
- d) Harlan
- 106. Which among the following scientists made pioneering contributions to wheat cytogenetics?
- a) C.R. Burnham
- b) B. McClintock
- c) Sanjay Rajaram
- d) E.R. Sears

- 107. The breakage-fusion-bridge cycle of chromosomes was first elegantly demonstrated by
- a) T.H. Morgan
- b) E.R. Sears
- c) C.R. Burnham
- d) B. McClintock
- 108. A *Drosophila* female with genotype AaBBCcddEeff when mated with a male of same genotype, how many different genotypes are expected in the progeny?
- a) 6
- b) 9
- c) 12
- d) 27
- 109. Lampbrush chromosomes become distinctively visible at which stage of meiosis in the oocytes?
- a) Leptotene
- b) Zygotene
- c) Diplotene
- d) Diakinesis
- 110. The pentose sugar molecule found in the genetic material of ØX174 is
- a) Ribose sugar
- b) Deoxyribose sugar
- c) Dideoxyribose sugar
- d) Both b) and c)
- 111. In a 3-line hybrid rice production system, which line is used as female parent?
- a) A line
- b) B line
- c) R line
- d) Mixture of B and R line
- 112. Which of the following is not associated with prokaryotic genome?
- a) Exon
- b) Intron
- c) Promoter
- d) TATA box
- 113. Which one of the following is an exclusive feature of the 'enhancer' elements?
- a) Enhancing and repressing gene expression
- b) Located upstream of the gene to be transcribed
- c) Orientation non-specificity
- d) Binding site for the DNA-binding proteins
- 114. Holandric gene is located on
- a) Autosome
- b) X chromosome
- c) Y chromosome
- d) Mitochondrial genome

- 115. 'Fairchild's Mule' is an
- a) Animal
- b) Insect
- c) Interspecific hybrid plant
- d) Invertebrate
- 116. In the absence of dominance or presence of codominance, which of the following formula can be used to predict the phenotypic classes in F<sub>2</sub>, where 'n' represents number of segregating genes?
- a) 2<sup>n</sup>
- b) 3<sup>n</sup>
- c) 4<sup>n</sup>
- d)  $2^{n+1}$
- 117. In a diploid rice plant, the number of chromosome in the endosperm cell would be
- a) 12
- b) 24
- c) 36
- d) 48
- 118. The amino acid having only one genetic code (codon) is
- a) Phenylalanine (Phe)
- b) Proline (Pro)
- c) Isoleucine (IIe)
- d) Tryptophan (Trp)
- 119. The DNA extracted from an F<sub>2</sub> population was amplified using a polymorphic SSR marker in a PCR. The amplified products were size separated on an agarose gel through gel electrophoresis and documented. How many different banding patterns can be expected on the gel?
- a) 1
- b) 2
- c) 3 d) 4
- 120. Which among the following has the highest power in identifying QTLs and estimating
- their effects?
  a) Single Marker Analysis
- b) Simple Interval Mapping
- c) Composite Interval Mapping
- d) Complex Interval Mapping
- 121. Which of the following is incorrect?
- a) SSRs are genetically codominant
- b) SSRs are hypervariable
- c) SSRs show multilocus profiles
- d) SSRs have abundant distribution in the genome

- 122. Northern blot hybridization is used for analysis of
- a) DNA
- b) RNA
- c) Proteins
- d) cDNA
- 123. 'Cry1' proteins are insecticidal to
- a) Lepidoptera
- b) Hymenoptera
- c) Diptera
- d) All of the above
- 124. Regarding sex determination in *Drosophila*, which of the following statements is not true?
- Y chromosome is primarily responsible for maleness
- b) Males have one X and one Y chromosome
- c) Ratio of sex chromosomes to autosomes determines the sex
- d) Females have two X chromosomes
- 125. Which among the following statements regarding a variety developed through MAS is correct? The MAS-derived variety will be
- Completely different from the recurrent parent variety
- Same as the recurrent parent variety except one or two traits
- c) Exactly similar to the donor parent only
- Intermediate between the recurrent and the donor parents
- 126. Protogyny is prevalent in
- a) Rice
- b) Wheat
- c) Soybean
- d) Mungbean
- 127. Linkage is apparent when recombination frequency (RF) is
- a) More than 50%
- b) Less than 50%
- c) Equal to 50%
- d) Both a) and b)
- 128. The unit of coefficient of variation (CV) is
- a) mm
- b) gm
- c) um
- d) None of the above
- 129. The basis of polymorphism detected by SSR marker is the
- Number of restriction sites in a particular sequence
- b) Number of repeats of a particular sequence
- c) Source (nuclear or mitochondrial) of the genome
- d) Size (large or small) of the genome

- 130. Pea (*Pisum sativum* L.) can have how many primary trisomics?
- a) 7
- b) 14
- c) 21
- d) 28

Matching type questions (No. 131 to 140); all questions carry equal marks. Choose the correct answer (a, b, c, d or e) for each sub-question (i, ii, iii, iv and v) and enter your choice in the circle (by shading with a pencil) on the OMR - answer sheet as per the instructions given on the answer sheet.

#### 131.

i) RFLP

- a) Kary Mullis
- ii) RAPD iii) AFLP
- b) Michelmore et al. (1991) c) Botstein et al. (1980)
- iv) PCR
- d) Vos et al. (1995)
- v) BSA
- e) Williams et al. (1990)

### 132.

i) Parthenocarpy

- a) Pearlmilletb) Rapeseed
- ii) Protandry
- c) Banana
- iii) Protogynyiv) Self-incompatibility
- d) Oat
- v) Cleistogamy
- e) Maize

### 133.

- i) Gene
- a) Waldeyer
- ii) Mutation
- b) Robert Hooke
- iii) Genetics
- c) H.J. Muller
- iv) Chromosome
- d) W. Johannsen

v) Cell

e) William Bateson

### 134.

- i) Gene-for-gene hypothesis
- a) Charles Yanofsky
- ii) Colinearity of gene and polypeptide
- b) Davenport
- iii) Chromosomal theory of inheritance
- c) Flor
- iv) Dominance hypothesis of heterosis
- d) G.H. Shull
- v) Single cross hybrid maize
- u) G.H. Shull
- v) Single cross hybrid maize
- e) Walter S. Sutton

## 135.

- i) Pure line theory
- a) Wr-Vr graph
- ii) Stability analysis
- b) Kempthorne (1957)
- iii) Diallele analysisiv) Path coefficient analysis
- c) Mahalanobis (1936) d) Johannsen (1903)
- v) D<sup>2</sup> analysis
- e) Eberhert and Russell (1966)

136.

i) Sir Archibald Garrod

- ii) Beadle and Tatum
- iii) Recombination between nucleotide pair
- iv) Deletion mapping

a) Charles Yanofsky

- b) H.G. Khorana
- c) Human metabolic diseases
- d) One-gene-oneenzyme
- v) Deciphering genetic code e) Seymour Benzer

137.

i) Annealing

a) Binding site of the polymerase enzyme

ii) Denaturation

b) Changing the genetic message in mRNA through deletion or addition of nucleotide

iii) Promoter

c) Binding of the primers to the template DNA

iv) RNA splicing

d) Separation of DNA strands

v) RNA editing

e) Removal of intron from premRNA

138.

i)  $SE_d \times 't'$ 

a) Mean variable / Standard deviation

ii) (Variance)<sup>1/2</sup>

b) Percent Standard deviation of means Standard deviation

iii) Coefficient of variance c) (CV)

iv) Normalization

d) Critical difference (CD)

v) Standard error (SE)

e) Standard deviation (SD)

of sample means

139.

i) Rice

ii) Soybean

iii) Potato

iv) Bread wheat v) Watermelon (seedless) a) Hexaploid

b) Tetraploid

c) Paleopolyploid

d) Triploid

e) Diploid

140.

i) Canola

iii) Barnase

v) Roundup-Ready

ii) Flavr-Savr

a) Pollen sterility

b) Insect resistance

c) Herbicide tolerant soybean

iv) Cry1Ac d) Long shelf-life tomato

e) Double-zero (00) rape

seed

Short questions (No. 141 to 146); each question carries FIVE marks. Write answers, including computation / mathematical calculations if any, in the space provided for each question on the question paper itself.

141. What do you mean by heterosis? Explain its components and show how the components can be manipulated to maximize heterosis?

142. Differentiate between multilines and pyramided lines, and write their merits and demerits.

143. Describe the role of markers in foreground and background selection in marker-assisted backcross breeding (MABB). List out the advantages of MABB over conventional backcross breeding.

144. What do you mean by 'reverse genetics'? How it differs from 'forward genetics'? Mention a few tools/techniques used in reverse genetics and their applications.

145. The genome of a crop (e.g. rice) has been sequenced. Explain, how the sequence information can be used for genetic improvement of the crop?

146. What is an euploidy? How does it differ from euploidy? Describe with example, how monosomics can be used in locating a dominant gene to a chromosome?